

p -Modular TQFT's and Torsion

THOMAS KERLER

THE OHIO STATE UNIVERSITY
COLUMBUS, OH, USA

September 6th, 2001

presented in Seminar/Workshop on

“Invariants of knots and 3-manifolds”

at the

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES

KYOTO, JAPAN

Non-Perturbative Quantum Invariants:

Let $p \geq 5$ be a prime, ζ_p primitive p -th root of unity.

$\mathcal{C}_p$ semisimple category over $\mathbb{Z}[\zeta_p]$ given by
even $(SO(3))$ part of $\overline{U_{\zeta_p}(\mathfrak{sl}_2) - mod}$

Quantum-Invariants:

M compact, oriented 3-manifold with $\partial M = \emptyset$.

[Reshetikhin, Turaev] $\mathcal{V}_{\zeta_p}^{RT}(M) \in \mathbb{C}$ w/ $\mathcal{V}_p^{RT}(S^3) = 1$.

[Murakami, Masbaum, Roberts] $\mathcal{V}_{\zeta_p}^{RT}(M) \in \mathbb{Z}[\zeta_p]$.

[Ohtsuki] For $\beta_1(M) = 0$ have $\lambda_j^O(M) \in \mathbb{Z}[\frac{1}{2}, \dots, \frac{1}{2j+1}]$

$$\mathcal{V}_{\zeta_p}^{RT}(M) = \sum_{j=0}^{\frac{p-3}{2}} \lambda_j^O(M) (\zeta_p - 1)^j + \mathcal{O}((\zeta_p - 1)^{\frac{p-1}{2}}).$$

[Murakami] $\lambda_{Casson} = 6\lambda_1^O$

TQFT:

[Atiyah] TQFT = Functor $\mathcal{V} : \mathbf{Cob} \longrightarrow \mathbf{R} - mod$

[Reshetikhin, Turaev] Constructions $\mathcal{V}_{\zeta_p}^{RT}$ for $\mathbf{R} = \mathbb{C}$.

[Gilmer] For $\mathbf{R} = \mathbb{Z}[\zeta_p]$ for “grounded” cobordisms.


[K] For $\mathbf{R} = \mathbb{Z}[\zeta_5]$ “half-projective” TQFT. Basis.

Topological Quantum Field Theory:

Cobordisms: $\partial M = \partial^+ M \sqcup \partial^- M$ w/ homeom.

$\psi^\pm : \partial^\pm M \xrightarrow{\sim} \Sigma_\pm$ to model surfaces.

Homeom. class $[M, \psi^\pm] : \Sigma_- \longrightarrow \Sigma_+$ (+Framings).



Category Cob: Composition by “Gluing” over surfaces.

TQFT Functor: Assigns $\mathcal{V}(\Sigma) \in \mathbf{R} - mod$ and for $M : \Sigma_- \rightarrow \Sigma_+$ have $\mathcal{V}(M) \in Hom_{\mathbf{R}}(\mathcal{V}(\Sigma_-), \mathcal{V}(\Sigma_+))$.

[K] “Half-Projective”: $\mathcal{V}(N \circ M) = x^{\mu(N, M)} \mathcal{V}(N) \mathcal{V}(M)$

with $\mu(N, M) = rk(H_1(N \circ M) \xrightarrow{\delta} H_0(M \cap N)), x \in \mathbf{R}$

Ordinary Composition: $x = 1$.

Mapping Class Group Representation:

$$\begin{aligned} \Gamma_g = \pi_0(Homeo^+(\Sigma)) &\xrightarrow{\cong} Aut(\Sigma) \xrightarrow{\mathcal{V}} GL(\mathcal{V}(\Sigma_g)) \\ [\psi] &\mapsto [\Sigma \times [0, 1], \psi \sqcup id] \end{aligned}$$

Main Question:

What is the TQFT content of the λ_j^O ?

Expansions $\mathcal{V}_{\zeta_p}^{RT}(M) = \sum_j \mathcal{V}_{p,j}^{RT}(M) y^j$ for $M \in \mathbf{Cob}$

via $\mathbb{Z}[\zeta_p] \twoheadrightarrow \mathbb{F}_p[\zeta_p] \cong \mathbb{F}_p[y] / y^{p-1}$

w/ $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$ and $y = \zeta_p - 1$

Applications & Remarks:

- Computations via TQFT-techniques
(Heegaard splits, Mapping tori, Seifert spaces, ...)

- Extensions of λ_j^O to $\beta_1(M) > 0$.
(Reidemeister Torsion vs. Casson Invariant)

- Filtration $\mathcal{M}_{p,k} = \bigcap_{j=0}^k \ker \left(\mathcal{V}_{p,j}^{RT} \Big|_{\Gamma_g} \right)$ of *mapping*

class group $\Gamma_g \stackrel{?}{\Longleftrightarrow} \text{Other (f.t.) Filtrations.}$

- Dependence on p : $\dim(\mathcal{V}_{\zeta_p}^{RT}(\Sigma_g)) \sim p^{3(g-1)}$ if $g > 1$
 $\implies$ NO fixed universal TQFT !
- Even for FIXED p , operators $\mathcal{V}_{p,j}^{RT}(M)$ over $\mathbb{F}_p$ are
NOT liftable to $\mathbb{Z}[\dots]$ as TQFT's.

Some Results - Overview:

General primes $p \geq 5$:

- Construct Family of *irreducible* TQFT's over $\mathbb{F}_p$
 $\overline{\mathcal{V}}_p^{(k)} \leftarrow (\overline{\mathcal{U}}_p^{(k)})$ for p odd prime and $0 < k < p(-3)$.
- Resolutions $0 \leftarrow \overline{\mathcal{V}}_p^{(k)} \leftarrow \mathcal{V}_p^{(k)} \leftarrow \mathcal{V}_p^{(2p-k)} \leftarrow \dots$
into $\mathbb{Z}$ -liftable TQFT's $\mathcal{V}_p^{(j)} \leftarrow \mathcal{V}_{\mathbb{Z}}^{(j)}$.
- Image of Alexander Polynomial $\overline{\Delta}_{p,\chi}(M) \in \mathbb{F}_p[\zeta_p]$
from traces over $\overline{\mathcal{V}}_p^{(k)}$ of covering cobordism.
- Similar results for p -modular S_n representations.

Fibonacci Case $p = 5$:

- $\mathcal{V}_{\zeta_5}^{RT}$ = half-projective, with $\mathbf{x} = \zeta_5 - \zeta_5^{-1} \in \mathbb{Z}[\zeta_5]$.
- Identification: $\mathcal{V}_{5,0}^{RT} = \overline{\mathcal{U}}_5^{(1)}$
- $2\overline{\Delta}_{5,\chi}(M) = \det(\ker(\chi)) + 3y\mathcal{V}_{\zeta_5}^{RT}(M) + \mathcal{O}(y^3)$
- $cut(M) \leq \mathfrak{o}_5(M)$ (joint w/ Gilmer)
- $\lambda_{Casson}(M_{D(\psi(\mathcal{C}_k))}) = \langle \Omega_{\Lambda}, \mathbb{P}_k^{\psi} \Omega_{\Lambda} \rangle \pmod{5}$
 $\Omega_{\Lambda} \in \bigwedge^g H_1(\Sigma_g)$, $\mathcal{C}_k$ k -th bounding curve; $\psi \in \Gamma_g$;
 $D(\mathcal{C})$ Dehn twist @ $\mathcal{C}$; M_{γ} Heegaard glue; $\mathbb{P}_k^2 = \mathbb{P}_k$.
- $\lambda_j^O \pmod{5}$ are finite type, $\mathcal{V}_{0,5}^{RT} = \det(H_1(M))$, ...

The Alexander Polynomial:

$M = \text{compact, oriented, } \beta_1(M) > 0$
 $\chi : H_1(M) \twoheadrightarrow \mathbb{Z}$ epimorphism
 $\widetilde{M}_\chi$
 $\downarrow$
 M
cyclic covering.

$H_1(\widetilde{M}_\chi, \mathbb{Z})$ is $\mathbb{Z}[t, t^{-1}]$ -module: t by Deck transf.

Definition: $\Delta_\chi(M) \in \mathbb{Z}[t, t^{-1}]$ as follows

- Maximal such that $E_1(H_1(\widetilde{M}_\chi)) \subseteq \Delta_\chi(M)\mathbb{Z}[t, t^{-1}]$
- $\det(A(t))$ if $A(t)$ is $n \times n$ -present. matrix of $H_1(\widetilde{M}_\chi)$.
- Choose symmetric (by P.D.) $\implies$ Unique up to sign.

Thm [Milnor, Turaev] Let $r_\chi(M)$ Reidemeister Torsion:

If $\partial M \neq \emptyset$ then $r_\chi(M) = (t - 1)^{-1} \Delta_\chi(M)$

If $\partial M = \emptyset$ then $r_\chi(M) = (t - 1)^{-2} \Delta_\chi(M)$

Remarks:

- For knot K : $M_K = S^3 - N(K)$ and $H_1(M_K) = \mathbb{Z}$.
- Multivariable $\Delta(M) \in \mathbb{Z}[H_1^{\text{free}}(M)]$
- $\Delta_\chi(1) = \det(\ker(\chi))$ w/ $\det(H) = \begin{cases} |H| & \text{if } < \infty \\ 0 & \text{elsewise} \end{cases}$
- $\Delta_\chi = 0$ if $\chi : \pi_1(M) \twoheadrightarrow \mathbb{Z} * \mathbb{Z} \twoheadrightarrow \mathbb{Z}$.
- $\Delta_p(S^1 \times \Sigma_g) = (t + t^{-1} - 2)^g$... dependence on $\chi!$
- **Thm [Meng, Taubes, Hutchings, Lee, Turaev, ...]**
Seiberg-Witten Invariant = (refined) *Torsion*.

TQFT's from Jacobians:

$$\mathcal{V}_{\mathbb{Z}}^{FN}(\Sigma) = H^*\left(\text{Hom}(\pi_1(\Sigma), U(1))\right) \cong \bigwedge^* H_1(\Sigma)$$

Thm [Frohman, Nicas]: Extends to TQFT

$$\mathcal{V}_{\mathbb{Z}}^{FN} : \mathbf{Cob} \longrightarrow \mathbb{Z} - mod$$

- $\mathbb{Z}/2\mathbb{Z}$ -projectivity/framing-dependence.
- Half-projective with $\kappa = 0$.
- $\Gamma_g \twoheadrightarrow Sp(2g, \mathbb{Z})$ Representation.
- $\mathcal{V}_{\mathbb{Z}}^{FN}(M) = \det(H_1(M, \mathbb{Z}))$ for $\partial M = \emptyset$.

Let $\Sigma \subset M$ 2-sided, oriented surface, dual to $\chi \in H^1(M)$

Thm [F,N] Set $C_{\Sigma} = (M - N(\Sigma)) : \Sigma \longrightarrow \Sigma$ in **Cob**.

$$\Delta_{\chi}(M) = \sum_{j=0}^{2g} (-t)^{j-g} \text{trace} \left(\mathcal{V}_{\mathbb{Z}}^{FN}(C_{\Sigma}) \Big|_{\bigwedge^j H_1(\Sigma_g)} \right)$$

Thm [K] $\mathcal{V}_{\mathbb{Z}}^{FN} \cong \mathcal{V}_{\mathcal{N}}^{Hennings}$ w/ $\mathcal{N} = \mathbb{Z}/2\mathbb{Z} \ltimes \bigwedge^* \mathbb{R}^2$

$\implies$ **[K]** Extension of Alexander-Conway Calculus to
3-manifolds presented by Surgery.

Note:

- **[F,N]** TQFT's via IH_* of $PSU(n)$ -Moduli Spaces.
- **[Donaldson]** TQFT's via connections on non-trivial $SO(3)$ -bundle (Casson-type) and SW (voretz)-equations.

Hard Lefschetz Decomposition:

Jacobians are Kähler $\Rightarrow$ Inner product on $\bigwedge^* H_1(\Sigma)$.

Symplectic 2-form $\omega = \sum_i a_i \wedge b_i \in \bigwedge^2 H_1(\Sigma)$.

$\mathfrak{sl}(2, \mathbb{Z})$ -Action: E, F, H generators act on $\bigwedge^* H_1(\Sigma)$ as

$$Hx = (\deg(x) - g)x, \quad Ex = x \wedge \omega, \quad F = E^*.$$

Thm: $\mathfrak{sl}(2, \mathbb{Z}) \xleftrightarrow{\text{DUAL on } \mathcal{V}_{\mathbb{Z}}^{FN}(\Sigma_g)} Sp(2g, \mathbb{Z})$

$$\text{hence} \quad \bigwedge^* H_1(\Sigma_g) = \bigoplus_{k=1}^g V_k \otimes W_{g,k}$$

where V_k is $k - \dim$ irred. $\mathfrak{sl}_2$ -module.

$W_{g,k}$ is irred. $Sp(2g, \mathbb{Z})$ -module; h.w. $\varpi_{g,k} = \sum_{i=1}^{g-k+1} \epsilon_i$

Thm [K]: $\mathcal{V}_{\mathbb{Z}}^{FN} \cong \bigoplus_{k=1}^{\infty} V_k \otimes \mathcal{V}_{\mathbb{Z}}^{(k)}$

- $\mathcal{V}_{\mathbb{Z}}^{(k)}(\Sigma_g) = \ker(F) \cap \bigwedge^{g-k+1} H_1(\Sigma_g) \cong W_{g,k}$
- $\mathcal{V}_{\mathbb{Z}}^{(k)}$ is irred. TQFT and $\mathcal{V}_{\mathbb{Z}}^{(k)}(\Sigma_g) = 0$ for $k > g+1$.
- $\Delta_{\chi}(M) = \sum_{k=1}^g [k]_{-t} \text{trace}(\mathcal{V}_{\mathbb{Z}}^{(k)}(C_{\Sigma}))$

$$\text{where } [n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}.$$

$\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$ -Reductions & Reducibility:

Let $p \geq 3$ be prime and $\mathcal{V}_p^{(j)}$ the $\mathbb{F}_p$ -reduction of $\mathcal{V}_{\mathbb{Z}}^{(j)}$.

Inherits inner form $\langle \cdot, \cdot \rangle_p : \mathcal{V}_p^{(j)}(\Sigma) \otimes \mathcal{V}_p^{(j)}(\Sigma) \longrightarrow \mathbb{F}_p$

Note! Forms $\langle \cdot, \cdot \rangle_p$ degenerate $\Rightarrow$ the $\mathcal{V}_p^{(j)}$ are Reducible!

Example:

Let $v = E1 - p(a_1 \wedge b_1) = \omega - p(a_1 \wedge b_1) \in \bigwedge^2 H_1(\Sigma_p)$.

Have $Fv = p - p = 0$ so that $v \in \ker(F) = \mathcal{V}^{(p-1)}(\Sigma_p)$.

For arbitrary $w \in \mathcal{V}^{(p-1)}(\Sigma_p)$ find

$$\begin{aligned} \langle v, w \rangle_p &= \langle E1, w \rangle_p - \langle p(a_1 \wedge b_1), w \rangle_p = \\ &= \langle 1, Fw \rangle_p - p \langle (a_1 \wedge b_1), w \rangle_p = 0. \end{aligned}$$

Note also:

$$E : \mathcal{V}_{\mathbb{Z}}^{(p+1)}(\Sigma_p) \rightarrow \mathcal{V}_{\mathbb{Z}}^{(p-1)}(\Sigma_p) + p \bigwedge^2 H_1(\Sigma_p, \mathbb{Z})$$

$$\text{so } E : \mathcal{V}_p^{(p+1)}(\Sigma_p) \rightarrow \mathcal{V}_p^{(p-1)}(\Sigma_p) \quad \text{well defined!}$$

Thm [K]: For $j \equiv k \pmod{p}$ with $0 < k < p$ have

$$E^k : \mathcal{V}_p^{(j)} \longrightarrow \mathcal{V}_p^{(j-2k)} \quad \text{well defined.}$$

All $E^k \neq 0$ and $E^p = 0$ in p -Reduction.

$\Rightarrow$ **Complex:** For any k with $0 < k < p$ have

$$\begin{aligned} \mathcal{C}(p, k) \quad \dots &\longrightarrow \mathcal{V}_p^{(c_{i+1})} \longrightarrow \mathcal{V}_p^{(c_i)} \longrightarrow \dots \\ \dots &\rightarrow \mathcal{V}_p^{(2p+k)} \longrightarrow \mathcal{V}_p^{(2p-k)} \longrightarrow \mathcal{V}_p^{(k)} \longrightarrow \overline{\mathcal{V}}_p^{(k)} \rightarrow 0 \end{aligned}$$

$$\text{w/ } c_i = \begin{cases} ip + k & i \text{ even} \\ (i+1)p - k & i \text{ odd} \end{cases} \quad \& \quad \text{maps } E^k \text{ or } E^{p-k}.$$

Denote: $\overline{\mathcal{V}}_p^{(j)}$ the subquotient by $\langle \cdot, \cdot \rangle_p$ -Null space.

Resolutions via Symmetric groups:

Thm [K] The sequences $\mathcal{C}(p, k)$ are *exact*.

The $\overline{\mathcal{V}}_p^{(k)}$ are *irreducible* with *resolutions* $\mathcal{C}(p, k)$.

Proof: $Sp(2g, \mathbb{Z})$ -weight spaces $\mathcal{W}^{(j)}(g, \lambda) \subset \mathcal{V}^{(j)}(\Sigma_g)$

with $\lambda = \sum_i \lambda_i \epsilon_i$ with $\lambda_j \in \{-1, 0, 1\}$

Let $N(\lambda) = \{i : \lambda_i = 0\}$ and $n(\lambda) = |N(\lambda)|$.

Via *Weyl Group* $S_{n(\lambda)}$ acts on $\mathcal{W}^{(j)}(g, \lambda)$ by permut. $N(\lambda)$.

Lem: $\mathcal{W}_{\mathbb{Z}}^{(j)}(g, \lambda) \cong \text{Specht-Mod. } \mathcal{S}_{\mathbb{Z}}^{\tau_j} \text{ (} S_{n(\lambda)}\text{-module)}$

with 2-row

Young Diagram
($n(\lambda)$ boxes)

$$\tau_j = \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \end{array} \dots \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline \end{array} \dots \underbrace{\begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline \end{array}}_{j-1} \dots$$

$\implies$ **Complex** of p -modular S_n -reps.

$$\dots \rightarrow \mathcal{S}_p^{\tau_{c_i}} \rightarrow \dots \rightarrow \mathcal{S}_p^{\tau_{2p-k}} \rightarrow \mathcal{S}_p^{\tau_k} \rightarrow \mathcal{D}_p^{\tau_k} \rightarrow 0$$

Jordan Hölder (Composition) Series:

$$0 \subset M_0 \subset M_1 \subset \dots \subset M_{m-1} \subset M_m = \mathcal{S}_p^{\tau}$$

All $R_i = M_i / M_{i-1} \neq 0$ and irreducible

Write $R_0 | R_1 | \dots | R_{m-1} | R_m$ and $\mathcal{F}(\mathcal{S}_p^{\tau}) = \{R_i\}$.

Thm [James] The $\mathcal{D}_p^{\tau}$ are irreducible.

$$\mathcal{F}(\mathcal{S}_p^{\tau_j}) = \{\mathcal{D}_p^{\tau_{j+2l}} : l \geq 0 \text{ and } l \leq \frac{p}{2} j + 2l\}$$

All multiplicities one or zero.

... not enough ...

Exactness via Modular Ordering:

Partial Order on $\mathcal{F}(\mathcal{S}_p^\tau)$ (w/ mult=0,1) defined by


$$R \leq R' \quad \text{iff} \quad R_0 | \dots | R | \dots | R' | \dots | R_m \quad \forall \text{ JH-Series.}$$

Thm [Kleshchev, Sheth, '99]

Determine $\leq$ for 2-row Young diagrams τ and $p \geq 3$:

$(\mathcal{F}(\mathcal{S}_p^\tau), \geq)$ “inclusion structure”
w/ $\mathcal{D}_p^\tau$ maximal element.

Cor [K] Modular structure of maps in $\mathcal{C}(p, k)$



with $\mathcal{F}(Y_i) = \mathcal{F}(X_i)$.

$\implies$ EXACTNESS. ■


Thm [K] Image of Alexander Polynomial wrt. $t \mapsto \pm \zeta_p$:

$$\overline{\Delta}_{\chi, p}^\mp(M) = \sum_{k=1}^{p-1} [k]_{\pm \zeta_p} \text{trace}(\overline{\mathcal{V}}_p^{(k)}(C_\Sigma)) \in \mathbb{F}_p[\zeta_p].$$

Johnson-Morita Extension:

Torelli Group: $0 \rightarrow \mathcal{I}_g \rightarrow \Gamma_g \rightarrow Sp(2g, \mathbb{Z}) \rightarrow 0$

Boundary Group: $\mathcal{K}_g$ smallest normal containing



Thm [Johnson, Morita] $\exists$ Homomorphisms $(H=H_1(\Sigma_g, \mathbb{Z}))$

$$\begin{array}{ccccccc}
 0 & \rightarrow & \mathcal{K}_g & \rightarrow & \Gamma_g & \xrightarrow{\tilde{k}} & \frac{1}{2} \frac{\Delta^3 H}{H} \rtimes Sp(2g, \mathbb{Z}) \rightarrow 0 \\
 & & \parallel & & \cup & & \cup \\
 0 & \rightarrow & \mathcal{K}_g & \rightarrow & \mathcal{I}_g & \xrightarrow{\tau_2} & \frac{\Delta^3 H}{H} \rightarrow 0
 \end{array}$$

Let $g^* Jg = J$ for $g \in Sp(2g, \mathbb{Z})$ and let $x \in \bigwedge^3 H$.

Define $\mu(x) : \bigwedge^j H \rightarrow \bigwedge^{j-3} H$ by $\langle a, \mu(x)b \rangle = \langle (Jx) \wedge a, b \rangle$.

Lem [K] μ factors into $Sp(2g, \mathbb{Z})$ -covariant map

$$\mu : \frac{1}{2} \frac{\Delta^3 H}{\omega \wedge H} \rightarrow Hom(\overline{\mathcal{V}}_p^{(k)}(\Sigma_g), \overline{\mathcal{V}}_p^{(k+3)}(\Sigma_g))$$

Thm [K] For $0 < k < p - 3$ obtain $\Gamma_g / \mathcal{K}_g$ -module

$$\begin{array}{l}
 \overline{\mathcal{U}}_p^{(k)}(\Sigma_g) = \overline{\mathcal{V}}_p^{(k)}(\Sigma_g) \overset{\mu}{\oplus} \overline{\mathcal{V}}_p^{(k+3)}(\Sigma_g) \\
 \text{by matrix} \quad (x, g) \mapsto \begin{bmatrix} g & 0 \\ \mu(x)g & g \end{bmatrix}
 \end{array}$$

Summary of TQFT Constructions:

	$\mathcal{V}_{\mathbb{Z}}^{FN}$	Jacobian TQFT Fully reducible over $\mathbb{Z}$
$\mathfrak{sl}_2$ -Lefschetz Decomposition	$\Downarrow$	(<i>Permutation Modules</i>)
	$\mathcal{V}_{\mathbb{Z}}^{(k)}$	Lefschetz Component, irreducible over $\mathbb{Z}$
p -Reduction	$\Downarrow$	(<i>Specht Modules</i> $\mathcal{S}_{\mathbb{Z}}^{(j)}$)
	$\mathcal{V}_p^{(k)}$	Reducible over $\mathbb{F}_p$ with inner form.
Null space quotient	$\Downarrow$	(<i>p-Specht Modules</i> $\mathcal{S}_p^{(j)}$)
	$\overline{\overline{\mathcal{V}}}_p^{(k)}$	Irreducible over $\mathbb{F}_p$. Not $\mathbb{Z}$ -liftable, but resolutions in $\mathcal{V}_p^{(j)}$'s
Johnson-Morita Extension	$\Downarrow$	(<i>Simple S_n-modules</i> $\mathcal{D}_p^{\{j\}}$)
	$\overline{\overline{\mathcal{U}}}_p^{(k)}$	Indecomposable with two factors $\overline{\overline{\mathcal{V}}}_p^{(k)}$ and $\overline{\overline{\mathcal{V}}}_p^{(k+3)}$


The Fibonacci TQFT:

$\mathcal{C}_5$ has two “colors” (simple objects) , namely 1 and ρ
 with $\rho \otimes \rho = 1 \oplus \rho$ and $1 = \text{unit}$.

RT-Vectorspace : over $\mathbb{Q}[\zeta_5]$

$$\mathcal{V}_{\zeta_5}^{RT}(\Sigma_g) = \text{Inv}_{\mathcal{C}_5}(\mathbf{F}^{\otimes g}) \quad \text{with} \quad \mathbf{F} = 1 \oplus (\rho \otimes \rho).$$

Fork basis: Example $g = 10$ ($1 \rightarrow \text{dotted}$; $\rho \rightarrow \text{solid}$)



Minimal Modules:

$$\mathfrak{S}_p(\Sigma) = \mathbb{Z}[\zeta_p] - \text{span of } \{\mathcal{V}_{\zeta_p}^{RT}(M)\Omega : M : \emptyset \rightarrow \Sigma\}$$

New Basis: Set $x = \zeta_5 - \zeta_5^{-1}$ and $D = -(\zeta_5^2 + \zeta_5^{-2})$.

$$\text{assign:} \quad U = \bigcup \quad [\dots] = \text{---} \boxed{\text{---} \cdot \cdot \cdot \text{---}} \text{---}$$

$$Y = \bigvee \quad W = \frac{1}{x} \left(\bigcup - D \bigcup \right)$$

and rescale with $x^{\lfloor \frac{\# \text{ of } Y's}{2} \rfloor}$ obtain new

$\Rightarrow$ basis vectors $\langle\langle [Y, W, U, Y, Y][W][Y, U, Y][U] \rangle\rangle$

denote: $\mathcal{V}_{\zeta_5}^I(\Sigma)$ the $\mathbb{Z}[\zeta_5]$ -span of these.

Thm [K]

$\exists_1$ half-projective TQFT $\mathcal{V}_{\zeta_5}^I$ with parameter x and

vectors spaces $\mathcal{V}_{\zeta_5}^I(\Sigma) = \mathfrak{S}_5(\Sigma)$

Proof: Explicit computation of **Cob**-generators.

Identification of $\mathbb{F}_5$ Reductions:

Denote: $\mathcal{V}_{0,5}^I$ the $\mathbb{Z}[\zeta_5] \rightarrow \mathbb{F}_5$ reduction of $\mathcal{V}_{\zeta_5}^I$.

Recall: With $0 < k = 1 < p - 3 = 2$ have construction

$$\begin{aligned} \overline{\mathcal{U}}_5^{(1)}(\Sigma_g) &= \overline{\mathcal{V}}_5^{(1)}(\Sigma_g) \overset{\mu}{\oplus} \overline{\mathcal{V}}_5^{(4)}(\Sigma_g) \\ \dots \quad \text{guess what} \quad \dots \end{aligned}$$

Thm [K] $\overline{\mathcal{U}}_5^{(1)} \cong \mathcal{V}_{0,5}^I$ (over $\mathbb{F}_5$).

Elements of Proof:

Middle dim in $\bigwedge^* \langle a_i, b_i \rangle$: $a_i \mapsto U$ and $b_i \mapsto W$.

$$\overline{\mathcal{V}}^{(1)} \cong \text{even } \# \text{ of } Y\text{'s} \quad \overline{\mathcal{V}}^{(4)} \cong \text{odd} \# \text{ of } Y\text{'s}$$

$$\text{Map} \begin{bmatrix} i \\ j \end{bmatrix} \leftrightarrow a_i \wedge b_i - a_j \wedge b_j$$

to Y -pair in i -th and j -th position



Resolve via T.L.-Recoupling theory into fork-tree gadget.

Dimensions: *Fibonacci* #'s : $f_{n+1} = f_n + f_{n-1}$
(with $f_0 = 0$, $f_1 = 1$, $f_2 = 1$, $f_3 = 2$, $\dots$)

For g even $\dim(\mathcal{V}_{0,5}^I(\Sigma_g)) = 5^{\frac{g}{2}} f_{g-1} \dots$

$\implies$ From Resolution of S_n -[Ryba]-reps. $\implies$

$$\begin{aligned} f_{2r} &= c(2r, r-1) - c(2r, r-3) + c(2r, r-6) - c(2r, r-8) + \\ &\quad + c(2r, r-11) - c(2r, r-13) + c(2r, r-16) - \dots \end{aligned}$$

$$\text{Catalan } \# \text{'s: } c(n, j) = \binom{n}{j} - \binom{n}{j-1}.$$

Applications of Fibonacci Identification:

- $2\overline{\overline{\Delta}}_{\chi,5}^+(M) = \dim(\ker(\chi)) - \mathfrak{x}\mathcal{V}_{\zeta_5}^{RT}(M) + \mathcal{O}(\mathfrak{x}^3).$

follows from character formula for $\overline{\overline{\Delta}}_{\chi,p}^\pm(M)$, and $\mathcal{V}^{RT}(M) = \mathfrak{x} \cdot \text{trace}(\mathcal{V}^{RT}(C_\Sigma))$ and Thm.

Note $\Rightarrow$ that if $\beta_1(M) > 1$ have

Thm[K] $\overline{\overline{\Delta}}_{\chi,5}^+(M) / \mathfrak{x}^3$ is independent of χ !!

$\Rightarrow$ Constraints on 3-manifold groups.

- If $\partial M = \emptyset$ then $\mathcal{V}_{\zeta_5}^{RT}(M) = \det(H_1(M)) + \mathcal{O}(\mathfrak{x}).$
(= Invariant of $\overline{\overline{\mathcal{V}}}^{(1)}$ and vacua lie in $\overline{\overline{\mathcal{V}}}^{(1)}$)

- The $\lambda_j^O \bmod 5$ are of finite type (known [Krick]).

Different Proof: Pick element in augmentation ideal $u \in$

II_g . Represented by $\mathcal{V}_{\zeta_5}^I$ in $\begin{bmatrix} 0 & 0 \\ \mu(u) & 0 \end{bmatrix} + \mathfrak{x} \cdot \text{End}.$

Thus $u^2 \in \mathfrak{x} \cdot \text{End}$ and $u^{2m} \in \mathfrak{x}^m \cdot \text{End}$. Also $v \in \mathcal{K}_g$ is represented in $\mathfrak{x} \cdot \text{End}$ so that $v^m \in \mathfrak{x}^m \cdot \text{End}$.

Hence $\lambda_{m-1}^O = 0$ for splittings in $(II_g)^{2m}$ and $(IK_g)^m$
 $\Rightarrow$ finite type by [Garoufalidis, Levine].

- Have $\mathcal{V}_{\zeta_5}^I(D(\mathcal{C}_k)) = \mathbb{I} + \mathfrak{x}\zeta_5^n \mathbb{P}_k$ with $\mathbb{P}_k^2 = \mathbb{P}_k.$

From $D(\psi(\mathcal{C}_k)) = \psi \circ D(\mathcal{C}_k) \circ \psi^{-1}$ we find

$$\mathcal{V}_{\zeta_5}^{RT}(D(\psi(\mathcal{C}_k))) = \mathbb{I} + \mathfrak{x}\overline{\overline{\mathcal{U}}}(\psi)\mathbb{P}_k\overline{\overline{\mathcal{U}}}(\psi)^{-1} + \mathcal{O}(\mathfrak{x}^2)$$

$$[\text{Murakami}] \Rightarrow \lambda_{\text{Casson}} = 2\langle \Omega, \overline{\overline{\mathcal{U}}}(\psi)\mathbb{P}_k\overline{\overline{\mathcal{U}}}(\psi)^{-1}\Omega \rangle$$

Cut-Numbers & Quantum-Orders:

Define: $cut(M)$ for closed, connected M by

- Largest $k \in \mathbb{N} \cup \{0\}$ s/th $\exists$ 2-sided $\Sigma \subset M$
with $\pi_0(\Sigma) = k$ but $\pi_0(M - \Sigma) = 1$.
- Largest $m \in \mathbb{N} \cup \{0\}$ s/th $\exists$ epimorphism
 $\pi_1(M) \twoheadrightarrow F_m \cong \underbrace{\mathbb{Z} * \dots * \mathbb{Z}}_m$ onto free group.

Thm [?] Two definitions are equivalent.

Note: $\beta_1(M) \geq cut(M)$ and
 $\beta_1(M) \neq 0 \Leftrightarrow cut(M) \neq 0$


Cor [K] Suppose $\mathcal{V}$ is half-projective w/ parameter x
Then $\mathcal{V}(M) \in x^{cut(M)} \cdot \mathbb{R}$.

Proof: Present

$M = A \circ B$

with

$\mu(A, B) = cut(M)$



Definition: Let $\mathfrak{o}_p(M)$ largest integer k such that

$$\mathcal{V}_{\zeta_p}^{RT}(M) \in (\zeta_p - 1)^k \mathbb{Z}[\zeta_p]$$

Thm [Gilmer, K] $cut(M) \leq \mathfrak{o}_5(M)$.

Thm [Melvin, Cochran] $\frac{\beta_1(M)}{3} \leq \frac{\mathfrak{o}_p(M)}{\mathfrak{o}_p(S^1 \times S^2)}$.

Two Examples [with Gilmer]:

Example 1 (0-surgery along links):

Let $M(\mathcal{L})$ be 3-mfld obtained via 0-surgery along $\mathcal{L} \subset S^3$

Lem: If $\mathcal{L}$ is boundary then $cut(M(\mathcal{L})) = |\mathcal{L}|$.

Denote: $\mathcal{B} = \mathcal{C}_1 \cup \mathcal{C}_2 \cup \mathcal{C}_3$ Borromean Link.

$\mathcal{C}^*$ the Whitehead Double of a component $\mathcal{C}$.

$\mathcal{B}^* = \mathcal{C}_1^* \cup \mathcal{C}_2 \cup \mathcal{C}_3$, $\mathcal{B}^{**} = \mathcal{C}_1^* \cup \mathcal{C}_2^* \cup \mathcal{C}_3$,

Clearly, $\beta_1(M(\mathcal{B}^{**})) = 3$. Note, that $M(\mathcal{B}) = (S^1)^{\times 3}$
so that $\pi_1(M(\mathcal{B})) = \mathbb{Z}^3$ and hence $cut(M(\mathcal{B})) = 1$.

Moreover, $\mathcal{B}^{**}$ is boundary so that $cut(M(\mathcal{B}^{**})) = 3$.


What is $cut(M(\mathcal{B}^*))$?

Have two surfaces $\mathcal{R}_1$ and $\mathcal{R}_2$.

Compute $\mathfrak{o}_5(M(\mathcal{B}^*)) = 2$

$\Rightarrow$ Answer $cut = 2$!

Note also: $\mathfrak{o}_5(M(\mathcal{B})) = 1$ and
 $\mathfrak{o}_5(M(\mathcal{B}^{**})) = 3$.



Example 2 (Mapping Tori):

For $j = 1, \dots, g-1$ let $\mathcal{A}_j \subset \Sigma_g$ be homologically independent, mutually non-intersecting curves, and $D_j = D(\mathcal{A}_j)$ the corresponding Dehn twists.

Let $M(n_1, \dots, n_{g-1}) \cong \Sigma_g \times [0, 1] / (x, 0) \sim (\psi(x), 1)$
be the mapping torus with $\psi = D_1^{n_1} \circ \dots \circ D_{g-1}^{n_{g-1}}$.

Prop: If all $n_j \not\equiv 0 \pmod{5}$ and $g \not\equiv 0 \pmod{5}$,
then $cut(M(n_1, \dots, n_{g-1})) = g$

Open Problems:

- Find $\mathbb{Z}[\zeta_p]$ -basis for $\mathcal{V}_{\zeta_p}^{RT}$ for $p \geq 7$.
 [Gilmer] Basis for $g = 1$ and *all* p .
 (? w/ [Masbaum] for $g = 2$?)
- Show they're half-projective w/ $x = (\zeta_p - \zeta_p^{-1})^{\frac{p-3}{2}}$.
 $\implies cut(M) \leq \frac{o_p(M)}{o_p(S^1 \times S^2)}.$
- Characterize $\xi(M) = \frac{\beta_1(M)}{cut(M)}$! Is it bounded?
 When is $\xi(M) \geq 3$? (Answers by [Sikora])
- Identify $\mathbb{F}_p$ -reduction $\mathcal{V}_{0,p}^I$ with $\overline{\mathcal{V}}_p^{(k)}$ and $\overline{\mathcal{U}}_p^{(k)}$.
NoGo Thm: For $p \geq 7$ cannot be linear relation.
Proof: by large g -asymptotics
 $dim(\overline{\mathcal{V}}_p^{(k)}(\Sigma_g)) \sim f_p^g$ and $dim(\mathcal{V}_{\zeta_p}^{RT}(\Sigma_g)) \sim F_p^g$.
 Have $f_5 = F_5$ but, e.g., $F_7 = 2f_7^2 - 7f_7 + 7$.
 Suggests a form $\mathcal{V}_{0,7}^I \subset \mathbb{F}_2 \otimes \overline{\mathcal{V}}_7^{(\#)} \otimes \overline{\mathcal{V}}_7^{(\#)}.$
- Is there TQFT for JM-extensions $\overline{\mathcal{U}}_p^{(k)}$? (By deformations of Jacobian-TQFT or $\mathbb{F}_2 \ltimes \bigwedge^* \mathbb{R}^2$ -theory?)
- What is structure of $\mathcal{V}_{j,p}^I$ for $j \geq 1$?
 $\implies$ Deeper structure of Mapping Class Group ...

- Use to find cohomology and characteristic classes of Mapping Class Group.
- For $\beta_1(M) > 0$ is $\mathcal{V}_{1,p}^I$ given by $\overline{\overline{\Delta}}_\chi(M)$. How?!
Relation to Frohman-Nicas- $PSU(n)$ -Knot Invariants?
- Which parts of $\overline{\overline{\Delta}}_{\chi,p}(M)$ independent of χ ?
Interpretation by $\mathbb{F}_p$ -covering, signature defects, etc.?
- What families of algebraic groups can be excluded from this as 3-manifold groups ?
- See $\mathcal{V}_{j,p}^I$ as “higher torsion” - invariants ?
 . . . as “ $\beta_1 > 0$ ”-extensions of λ_j^O ?
- Relations to 3D gauge theory (Seiberg Witten, Donaldson)
- Reexpress formulae for $\lambda_{Casson} \bmod 5$ for Heegaard splits with $g \geq 2$ in $Sp(2g, \mathbb{Z})$ -invariant-theoretical terms.
Compare to Morita’s formulae!
For $g = 2$ know $\mathbb{P}_1$ explicitly
 $\Rightarrow$ precisely reproduce Morita formula.
(looks a-priori different for $g > 2$)
- Same for general p .
- Heegaard formulae also for $\lambda_j^O \bmod p$ for $j > 1$ and general p .